



10 Points in Dimension 4 not Projectively Equivalent to the Vertices of a Convex Polytope

DAVID FORGE, MICHEL LAS VERGNAS[†] AND PETER SCHUCHERT

Using oriented matroids, and with the help of a computer, we have found a set of 10 points in \mathbb{R}^4 not projectively equivalent to the vertices of a convex polytope. This result confirms a conjecture of Larman [6] in dimension 4.

© 2001 Academic Press

PROBLEM (McMullen [6]). Determine the largest integer $n = f(d)$ such that for any given n points in general position in \mathbb{R}^d there is an admissible projective transformation mapping these points onto the vertices of a convex polytope.

Here admissible means that none of the n points is sent to infinity by the projective transformation.

For dimension two and three the numbers $f(d)$ are known: $f(2) = 5$ and $f(3) = 7$. For $d \geq 2$, Larman has established in [6] the bounds $2d + 1 \leq f(d) \leq (d + 1)^2$, and conjectured that $f(d) = 2d + 1$. The upper bound has been improved to $f(d) \leq (d + 1)(d + 2)/2$ by Las Vergnas [7], as a corollary of Redei's theorem for tournaments. Recently, Ramírez Alfonsín [8] has proven the linear upper bound $f(d) \leq 5d/2 + 1$, by a construction using Lawrence oriented matroids (unions of rank 1 oriented matroids).

In the context of oriented matroids the problem can be conveniently restated in terms of hyperplanes. We refer the reader to [1] for information regarding oriented matroid theory. As easily seen, the oriented matroids of the images of a given configuration of points by admissible projective transformations are all the acyclic reorientations of the oriented matroid defined by the affine dependencies of the configuration. The dual of a configuration of points is an arrangement of hyperplanes, and the regions defined by this arrangement are in 1–1 correspondence with the acyclic reorientations of the oriented matroid. We say that a region which meets all hyperplanes in dimension $d - 1$ is *complete*. It is almost immediate to verify that a region is complete if and only if all corresponding admissible projective transformations maps the given n points onto the set of vertices of convex polytopes (note that these convex polytopes necessarily have the same oriented matroid).

Hence the McMullen problem is equivalent to: *determine the largest integer $n = f(d)$ such that any arrangement of n hyperplanes in general position in \mathbb{R}^d contains a complete region.*

The same problem for general oriented matroids has been considered by Cordovil and Da Silva [4]: *determine the largest integer $n = g(r)$ such that any uniform rank r oriented matroid M with n elements has a complete region.* A *region* (or *tope*) of an oriented matroid is a region determined by the pseudohyperplanes of its topological representation. The regions of an oriented matroid are in 1–1 correspondence with its maximal covectors, and a region is complete if and only if changing the sign of any element in the corresponding maximal covector produces another maximal covector. Obviously $g(r) \leq f(r + 1)$. Cordovil and Da Silva have shown in [4] that $2r - 1 \leq g(r)$, generalizing Larman's lower bound.

In this paper, we construct uniform rank 5 oriented matroids on 10 elements without complete region, hence, $g(5) = 9$. One of these oriented matroids has a realization in \mathbb{R}^4 , hence $f(4) = 9$.

[†]C.N.R.S., Paris.

As a preliminary step for the rank 5 case, using a computer, we have gone through the complete list of all 2628 reorientation classes of uniform rank 4 oriented matroids on eight elements, as per the work of Bokowski and Richter-Gebert [2].

PROPOSITION 1. *There are precisely 114 non-isomorphic reorientation classes of uniform rank 4 oriented matroids on eight elements without complete region. One such reorientation class has only mutants without complete region. Two of them are not realizable.*

The unique realizable uniform rank 4 oriented matroid on eight elements without complete region, such that all its mutants are also without complete region, has the following base signature (or chirotope):

+++++ - - - - - + - - - - - - - - - + + + +
- + - - - - - + - - - - - - - - - + + + + +

THEOREM 2. *There is a set of 10 points of \mathbb{R}^4 in general position such that:*

- *there is no admissible projective transformation mapping these points onto the vertices of a convex polytope, or, equivalently,*
- *the corresponding uniform oriented matroid has no complete region.*

The theorem means that $f(4) = g(5) = 9$.

PROOF. Using a computer, it can be checked that the oriented matroid of affine dependencies of the following 10 points of \mathbb{R}^4 has no complete region.

| | | | | |
|----|--------|--------|---------|-------|
| 1 | 0.7702 | 0.2217 | -6.3645 | 0 |
| 2 | 0.7426 | 0.2284 | -6.3977 | 0 |
| 3 | 0.6 | 1.01 | -5.44 | 0 |
| 4 | 1.75 | 7.07 | -0.45 | 0 |
| 5 | -2 | 2 | 2 | 1 |
| 6 | 2 | -2 | 2 | 1 |
| 7 | 2 | 2 | -2 | 1 |
| 8 | -2 | -2 | -2 | 1 |
| 9 | -2.44 | -2.13 | 1.4 | 1.71 |
| 10 | 0.35 | 1.77 | -0.38 | 1.011 |

□

The signature of the 252 bases of this uniform rank 5 oriented matroid on 10 elements is:

+++++ - - - - - + - - - - - + + + - - - - - - - - - + - - + + - - - - - + - - + +
+ - - - - + - - - - - + - + + - + - - - - - + + + + - + - + + + - + - + + + +
- - - + + + + + + - - - - - + + + + - - + + + + + + - - - + + + + - - - + -
+ + + + + - - - - - + + + + - + - + + - - + + + + + - - - + + + + + + - - -
- + + + + - - - + + + + + + - - - + + + - - - + - + + + - - - + + + + -
- - + + + + + + - - - + + + + - - - + - + + + - - - + + + + - - - + - - + - + -

Its face lattice is that of a stacked 4-cross-polytope with 19 facets (we recall that a *stacked* polytope is obtained by the addition of new vertices building shallow pyramids over facets):

1234 1238 1247 1278 1346 1368 1467 1678 2345 2358
2457 2578 3456 3568 4567 5679 5689 5789 6789

The point 9 is stacked on the 4-cross-polytope by the vertices 1, ..., 8 and the point 10 lies inside the convex hull of the points 5, 6, 7, 8 and 9. The vertices 5, 6, 7 and 8 form a

regular tetrahedron. The computer program provides the number of regions adjacent to each of the 256 regions of the oriented matroid: there are 16 with five neighbours, 57 with six neighbours, 72 with seven neighbours, 65 with eight neighbours, 46 with nine neighbours and 0 with 10 neighbours.

We now explain how we arrived to our example. Since a list of all reorientation classes of uniform rank 5 oriented matroids on 10 elements does not exist we cannot use exhaustion as in the rank 4 case.

We start with the list of 135 reorientation classes of uniform rank 5 oriented matroids on eight elements [2, 3]. From this list we can generate the 3501 non-isomorphic matroid polytopes of rank 5 with eight vertices. The face lattices of these matroid polytopes are the 37 3-spheres with eight vertices described by Grünbaum and Sreedharan [5].

For any such matroid polytope P and any disjoint pair of facets f_1, f_2 of its face lattice we generate a partial uniform rank 5 oriented matroid M on 10 elements as follows. The face lattice of M is a stacked 3-sphere where the vertex 9 is stacked on f_1 in the 3-sphere P . The element 10 of M is an interior element with a special relationship to some of its combinatorial hyperplanes. The facets f_1 and f_2 each have four elements. Let H_{31} resp. H_{13} be a combinatorial hyperplane with three elements of f_1 and 1 of f_2 resp. three elements of f_2 and one of f_1 . Then the element 10 lies on the same side of H_{31} as the element of $f_1 \setminus H_{31}$ and on the same side of H_{13} as the element of $f_2 \setminus H_{13}$. In this way we can construct 18 872 partial oriented matroids. Starting from these partial oriented matroids, we generate 1112 uniform rank 5 oriented matroids on 10 elements without complete region. They lie in 414 reorientation classes. If we build the mutants of these oriented matroids, we come up to 465 non-isomorphic reorientation classes of oriented matroids without complete region. None of them has all its mutants without complete region.

THEOREM 3. *There are at least 465 non-isomorphic reorientation classes of uniform rank 5 oriented matroids on 10 elements without complete region.*

ACKNOWLEDGEMENT

The third author (PS) wishes to thank the Equipe Combinatoire for the nice atmosphere during his visit to Paris. He also acknowledges partial support from the European HCM-Network DIMANET ERBCHRXCT 94 0429.

REFERENCES

1. A. Björner, M. Las Vergnas, B. Sturmfels, N. White and G. Ziegler, *Oriented Matroids*, Cambridge University Press, 1993.
2. J. Bokowski, Oriented matroids, in: *Handbook of Convex Geometry*, Chapter 2.5, P. M. Gruber and J. M. Wills (eds), North-Holland, 1993.
3. J. Bokowski, Finite point sets and oriented matroids, in: *Learning and Geometry*, W. Kucker and C. H. Smith (eds), Birkhäuser, 1996.
4. R. Cordovil and I. P. Silva, A problem of McMullen on the projective equivalences of polytopes, *Europ. J. Combinatorics*, **6** (1985), 157–161.
5. B. Grünbaum and V. Sreedharan, An enumeration of simplicial 4-polytopes with 8 vertices, *J. Comb. Theory*, **2** (1967), 437–465.
6. D. G. Larman, On sets projectively equivalent to the vertices of a convex polytope, *Bull. London Math. Soc.*, **4** (1972), 6–12.

7. M. Las Vergnas, Hamilton paths in tournaments and a problem of McMullen on projective transformations in \mathbb{R}^d , *Bull. London Math. Soc.*, **18** (1986), 571–572.
8. J. L. Ramírez Alfonsín, Lawrence oriented matroids and a problem of McMullen on projective equivalence of polytopes, *Europ. J. Combinatorics*, **22** (2001), 723–731, doi: 10.1006/eujc.2000.0492.

Received 31 January 2000 in revised form 4 December 2000

DAVID FORGE AND MICHEL LAS VERGNAS

*Université Paris 6,
case 189 - Combinatoire, 4 Place Jussieu,
75005 Paris,
France*

E-mail: forge@ccr.jussieu.fr; mlv@ccr.jussieu.fr

AND

PETER SCHUCHERT

*Technische Hochschule Darmstadt,
Schloßgartenstr. 7,
D-64289 Darmstadt,
Germany*

E-mail: schucher@mathematik.th-darmstadt.de